
Jos Marn
**Preconditioners for nonsymmetric linear systems with
low-rank skew-symmetric part**

Institut de Matematica Multidisciplinar
Universitat Politecnica de Valencia
Cam de Vera
14
46022 Valencia
Spain
`jmarinma@imm.upv.es`
Juana Cerdn
Danny Guerrero
Jos Mas

In this work we study the iterative solution of nonsingular, nonsymmetric linear systems of n equations

$$Ax = b \tag{1}$$

where the skew-symmetric part of the coefficient matrix A can be approximated by a low-rank matrix. Consider $A = H + K$ where H and K are the symmetric and skew-symmetric parts of A , respectively. It is assumed that the skew-symmetric matrix can be written as $K = PQ^T + E$ for some full rank $P, Q \in \mathbb{R}^{n \times s}$ with $k \ll n$, and $\|E\| \ll 1$.

Different strategies have been proposed when the skew-symmetric part K has exactly rank $s \ll n$. In [1] it is presented a progressive GMRES method that allows for the short-term computation of an orthogonal Krylov subspace basis. As pointed out in [5], although the method is mathematically equivalent to full GMRES [6] in practice it may suffer from instabilities due to the loss of orthogonality between the vectors of the generated Krylov subspace basis. In the same paper, the authors propose a Schur complement method that also permits the application of short-term formulas. The method obtains an approximate solution by applying the MINRES method $s+1$ times. The authors also suggest that can be successfully applied as a preconditioner for GMRES for the problem considered in this paper.

We study a method based on the framework proposed in [4]. Assuming that the matrix $H + PQ^T$ is nonsingular, our approach computes an approximate LU factorization of the matrix

$$\begin{bmatrix} H & P \\ -Q^T & I \end{bmatrix} \tag{2}$$

with the Balanced Incomplete Factorization (BIF) algorithm [2, 3]. Interestingly, the matrix in (2) is similar to the one used in [5] to develop the Schur

complement method, but in this work it is used to update a previously computed preconditioner for the symmetric part H . Then, the factorization is used as a preconditioner for the GMRES method. The results of the numerical experiments for different problems will be presented.

Bibliography

- [1] Bernhard Beckermann and Lothar Reichel. The Arnoldi process and GMRES for nearly symmetric matrices. *SIAM J. Matrix Anal. Appl.*, 30(1):102–120, February 2008.
- [2] R. Bru, J. Marín, J. Mas, and M. Tũma. Balanced incomplete factorization. *SIAM J. Sci. Comput.*, 30(5):2302–2318, 2008.
- [3] R. Bru, J. Marín, J. Mas, and M. Tũma. Improved balanced incomplete factorization. *SIAM J. Matrix Anal. Appl.*, 31(5):2431–2452, 2010.
- [4] J. Cerdán, J. Marín, and J. Mas. Low-rank updates of balanced incomplete factorization preconditioners. *Submitted*.
- [5] Mark Embree, Josef A. Sifuentes, Kirk M. Soodhalter, Daniel B. Szyld, and Fei Xue. Short-term recurrence Krylov subspace methods for nearly hermitian matrices. *SIAM J. Matrix Anal. and Appl.*, 33-2:480–500, 2012.
- [6] Y. Saad and M. H. Schulz. GMRES: A generalized minimal residual algorithm for solving nonsymmetric linear systems. *SIAM Journal on Scientific and Statistical Computing*, 7:856–869, 1986.