
John Shadid
**Scalable Solution of Continuum Plasma Physics Models
by Approximate Block Factorization Preconditioners**

Sandia National Laboratories
Computational Mathematics Department
Sandia National Laboratories
PO Box 5800 MS 1321
Albuquerque
NM 87185 1321
`jnshadi@sandia.gov`
Edward Phillips
Eric Cyr
Roger Pawlowski

The mathematical basis for the continuum modeling of plasma physics systems is the solution of the governing partial differential equations (PDEs) describing conservation of mass, momentum, and energy, along with various forms of approximations to Maxwell's equations. This PDE system is non-self adjoint, strongly-coupled, highly-nonlinear, and characterized by physical phenomena that span a very large range of length- and time-scales. To enable accurate and stable approximation of these systems a range of spatial and temporal discretization methods are commonly employed. In the context of finite element spatial discretization methods these include mixed integration, stabilized methods and structure-preserving (physics compatible) approaches. For effective time integration for these systems some form of implicitness is required. Two well-structured approaches, of recent interest, are fully-implicit and implicit-explicit (IMEX) type methods. Clearly the desire to accommodate disparate spatial discretizations, and allow the flexible assignment of mechanisms as explicit or implicit operators, implies a wide variation in unknown coupling, ordering, the nonzero block structure, and the conditioning of the implicit system. These characteristics make the scalable and efficient iterative solution of these systems extremely challenging.

This talk considers the development and evaluation of approaches based on approximate block factorization (ABF) and physics-based preconditioning approaches. These methods reduce the coupled systems into a set of simplified systems to which multilevel methods are applied. A critical aspect of these methods is the development of approximate Schur complement operators that encode the critical cross-coupling physics of the system [1-3]. To demonstrate the flexibility and performance of these methods we consider application of these techniques to resistive MHD and multiphysics MHD models for challenging prototype systems. In this context robustness, efficiency, and the parallel

and algorithmic scaling of the preconditioning methods are discussed.

[1] E. C. Cyr, J. N. Shadid, R. S. Tuminaro, R. P. Pawlowski, and L. Chacon, A new approximate block factorization preconditioner for 2D incompressible (reduced) resistive mhd, SIAM Journal on Scientific Computing, 35:B701-B730, 2013

[2] E. G. Phillips, H. C. Elman, E. C. Cyr, J. N. Shadid, R.P. Pawlowski, A Block Preconditioner for an Exact Penalty Formulation for Stationary MHD, SIAM J. Sci. Comput. 36-6 (2014), pp. B930-B951

[3] E. C. Cyr, J. N. Shadid, and R. S. Tuminaro, Teko an abstract block preconditioning capability with concrete example applications to Navier-Stokes and resistive MHD, Accepted in SISC

*This work was supported by the DOE Office of Science Advanced Scientific Computing Research - Applied Math Research program at Sandia National Laboratories.