
Christoph Pflaum
**Discretization of Elliptic Differential Equations Using
Sparse Grids and Prewavelets**

Friedrich-Alexander-Universitt Erlangen-Nrnberg
Department Computer Science 10
Cauerstrae 11
D-91058 Erlangen
`christoph.pflaum@fau.de`
Rainer Hartmann

Sparse grids can be used to discretize second order elliptic differential equations on a d -dimensional cube. Using Galerkin discretization, we obtain a linear equation system with $O(N(\log N)^{d-1})$ unknowns. The corresponding discretization error is $O(N^{-1}(\log N)^{d-1})$ in the H^1 -norm. A major difficulty in using this sparse grid discretization is complexity of the related stiffness matrix. Consequently, only differential equations with constant coefficients could be efficiently discretized using sparse grids for $d > 2$. To reduce the complexity of the sparse grid discretization matrix, we apply pre-wavelets. This simplifies the implementation of the multigrid Q-cycle. Furthermore, we present a new sparse grid discretization for Helmholtz equation with a variable coefficient c . This discretization utilizes a semi-orthogonality property. The convergence rate of this discretization in H^1 -norm is $O(N^{-1}(\log N)^{d-1})$ for $d \leq 4$ and $O(N^{-(2/(d-2))})$ for $d \geq 5$.